

# DATASCI 347: Machine Learning

## Lecture 1: Preliminaries in machine learning

Ruoxuan Xiong

# Lecture plan

- Preliminaries in machine learning
  - Parametric vs. nonparametric methods
  - Training vs. test data
  - Training vs. test mean squared error (MSE)





# Supervised vs unsupervised learning

- **Supervised learning** (main focus of this course)
  - **Data:**  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ 
    - $X_i$ : features (predictors)
    - $Y_i$ : label/response
  - **Goal:** learn a function  $f$  so that  $\hat{Y} = f(X)$
  - **Examples:** regression, classification (e.g., linear/logistic regression, tree-based methods, neural network)
- **Unsupervised learning**
  - **Data:**  $X_1, X_2, \dots, X_n$
  - **Goal:** discover structure/patterns in  $X$  (no labels)
  - **Examples:** clustering, dimensionality reduction (PCA)



# Supervised machine learning: How do we estimate $f$ ?

- Supervised machine learning finds a **function  $f$**  that maps  $X$  to  $Y$
- We may first look at the scatterplot for the exploratory analysis




$$Y = \beta_0 + X \cdot \beta_1 + \varepsilon$$


# Supervised machine learning: How do we estimate $f$ ?

- Supervised machine learning finds a **function  $f$**  that maps  $X$  to  $Y$
- We may first look at the scatterplot for the exploratory analysis

Simulated Dataset 2



Simulated Dataset 2



$$Y = \beta_0 + X \cdot \beta_1 + X^2 \cdot \beta_2 + \varepsilon$$


# Parametric methods

- We assume that  $f$  takes a specific form indexed by *a fixed number of parameters*.  
For example,
  - $Y = \beta_0 + X \cdot \beta_1 + \varepsilon$
  - $Y = \beta_0 + X \cdot \beta_1 + X^2 \cdot \beta_2 + \varepsilon$
- We use the training data,  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ , to *fit* the parameters


# A more complicated case...

- From the scatterplot, which  $f$  should we choose?


Simulated Dataset 1



Simulated Dataset 1




Simulated Dataset 1



# Nonparametric methods


- We don't make any assumptions on the form of  $f$ , but we restrict how “rough” the function can be
  - For example,  $k$ -nearest neighbors (KNN)

Simulated Dataset 1




Roughest

Simulated Dataset 1





Simulated Dataset 1



Smoothest

# Parametric vs nonparametric methods

- **Parametric methods** are often simpler to interpret, but strongly rely on assumptions and can be less flexible to capture complex data patterns
- **Nonparametric methods** rely on fewer assumptions, are flexible and suitable for large datasets




# In practice, which model we should use?


- Linear model, quadratic model, nonparametric model — or something else?
- To compare models, we need an **evaluation metric**
- From regression analysis, one common choice is  $R^2$  (goodness of fit)
  - $$R^2 = 1 - \frac{\sum_i (Y_i - \hat{Y}_i)^2}{\sum_i (Y_i - \bar{Y})^2}$$
  - $\hat{Y}_i$  is the fitted  $Y_i$  and  $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$ 
    - In linear regression,  $\hat{Y}_i = \hat{\beta}_0 + X_i \cdot \hat{\beta}_1$
    - For a more general fitted function  $\hat{f}$  (e.g., quadratic),  $\hat{Y}_i = \hat{f}(X_i)$
  - **Interpretation of  $R^2$** : Fraction of the variance of  $Y_i$  captured by  $\hat{f}(X_i)$ . The larger the  $R^2$ , the better  $\hat{Y}_i$  fits  $Y_i$
  - *Quiz: Can  $R^2$  be less than zero? Can  $R^2$  be larger than one?*




# Example of $R^2$




$$R^2 = 0.138$$




$$R^2 = 0.166$$



$$R^2 = 0.305$$



$$R^2 = 0.191$$




$$R^2 = 0.158$$


# Mean-squared error (MSE)

- MSE and RMSE are commonly used to evaluate regression models
  - $\text{MSE} = \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{f}(X_i))^2$
  - $\text{MSE} \geq 0$
  - If  $\hat{f}(X_i)$  is very close to  $Y_i$  for all  $i$ , then MSE would be small
  - $R^2 = 1 - \frac{\sum_i (Y_i - \hat{Y}_i)^2}{\sum_i (Y_i - \bar{Y})^2}$  is the standardized version of MSE
  - Root Mean-Squared Error (RMSE) is  $\sqrt{\text{MSE}}$


# MSE and RMSE




MSE = 0.439  
RMSE = 0.663




MSE = 0.425  
RMSE = 0.652



MSE = 0.354  
RMSE = 0.595



MSE = 0.412  
RMSE = 0.642




MSE = 0.429  
RMSE = 0.655

# MSE, RMSE and $R^2$


- Given the training data,  $(X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n)$ , there is a one-to-one mapping between MSE, RMSE and  $R^2$
- When is each metric used?
- **MSE**: (a) used in **model training** because it is mathematically simpler and differentiable; (b) used in **theoretical analysis**
- **RMSE**: Used in **performance reporting** because it reflects **error in original data scale**
- **$R^2$** : A **scale-independent metric**, used when audience is familiar with “**percentage of variance explained**”




# One-to-one mapping between MSE, RMSE and $R^2$




MSE = 0.439  
RMSE = 0.663  
 $R^2 = 0.138$




MSE = 0.425  
RMSE = 0.652  
 $R^2 = 0.166$



MSE = 0.354  
RMSE = 0.595  
 $R^2 = 0.305$




MSE = 0.412  
RMSE = 0.642  
 $R^2 = 0.191$




MSE = 0.429  
RMSE = 0.655  
 $R^2 = 0.158$

# Scaling of MSE, RMSE and $R^2$

If we multiply  $x_i$  and  $y_i$  by 2




$$\begin{aligned} \text{MSE} &= 0.439 \\ \text{RMSE} &= 0.663 \\ R^2 &= 0.138 \end{aligned}$$




$$\begin{aligned} \text{MSE} &= 1.756 \\ \text{RMSE} &= 1.325 \\ R^2 &= 0.138 \end{aligned}$$

# Which model to use for prediction?


- Suppose we have  $m$  new units
  - Their predictors are  $X'_1, X'_2, \dots, X'_m$
  - We want to predict the outcome of these  $m$  units
  - *Quiz: Which model should we use? Shall we choose the one with minimum MSE?*




MSE = 0.439  
RMSE = 0.663  
 $R^2 = 0.138$




MSE = 0.425  
RMSE = 0.652  
 $R^2 = 0.166$



MSE = 0.354  
RMSE = 0.595  
 $R^2 = 0.305$




MSE = 0.412  
RMSE = 0.642  
 $R^2 = 0.191$




MSE = 0.429  
RMSE = 0.655  
 $R^2 = 0.158$

# Which model to use for prediction?


- Suppose we have  $m$  new units
  - Their predictors are  $X'_1, X'_2, \dots, X'_m$
  - The fitted outcomes are  $\hat{Y}'_1, \hat{Y}'_2, \dots, \hat{Y}'_m$
- Suppose we are clairvoyants, and know the true outcome  $Y'_1, Y'_2, \dots, Y'_m$ 
  - We can calculate  $\text{MSE} = \frac{1}{m} \sum_{i=1}^m (Y'_i - \hat{f}(X'_i))^2$




MSE = 0.533




MSE = 0.518



MSE = 0.564




MSE = 0.507



MSE = 0.524

# *A low training MSE does not imply a low test MSE...*

- This is the main challenge in machine learning




Training MSE = 0.439

Training MSE = 0.425

Training MSE = 0.354

Training MSE = 0.412

Training MSE = 0.429



Test MSE = 0.533

Test MSE = 0.518

Test MSE = 0.564


Test MSE = 0.507

Test MSE = 0.524



# MSE varies with model flexibility


More flexible model



Training MSE = 0.439

Training MSE = 0.425

Training MSE = 0.354



Test MSE = 0.533

Test MSE = 0.518

Test MSE = 0.564

